

IASSNS-HEP-98/32

hep-th/9804019

April 1998

Supergravity, the DBI Action and Black Hole Physics

S. P. de Alwis^{1 2}

School of Natural Sciences, Institute for Advanced Study, Princeton NJ 08540

Abstract

The assumptions behind the recently conjectured relation between gauge theory and supergravity are elaborated on. It is pointed out that the scaling limit that preserves supergravity solutions, gives the entire DBI action on the gauge theory side, but in the low energy limit the relation between the conformal field theory and Anti-de Sitter supergravity emerges. We also argue that recent work on these issues may help in understanding the physics of five (four) dimensional black hole with three (four) charges in the so-called dilute gas region.

¹e-mail: dealwis@sns.ias.edu, dealwis@gopika.colorado.edu

²On leave from Department of Physics, Box 390, University of Colorado, Boulder, CO 80309.

1 Introduction

Recent work has highlighted the correspondence between supergravity solutions for black holes (branes) and their thermodynamic properties, with microscopic properties of D-brane gauge theory (for recent reviews see [1]). One important case that has been studied in some detail has been the absorption of scalars from D3 branes [2] where it was shown that exact agreement exists between certain supergravity calculations and the gauge theory calculation. On the basis of these results it has been conjectured [3] that there is a “new type of duality” between gauge theory and gravity. In [4] and [5] this conjecture was given a precise form in terms of an ansatz relating the supergravity action at a classical solution to a certain boundary conformal field theory. To understand this connection better it would be necessary to formulate string theory in Ramond-Ramond backgrounds. However so far it has not been possible to find a conformal field theory in such backgrounds. The best one can do is to consider the operator formulation of string perturbation theory in which one can construct vertex operators for the RR field strengths.

In order to find the relation to gauge theory we need to allow the string world sheet to have boundaries which are attached to D-branes (see [6] for a review). The techniques for dealing with this situation were developed in [7] (for D9-branes) and were adapted for use in the general D-brane situation in [8],[9],[10]. These arguments may be summarized as follows. Let the sum of vertex operators for the fluctuations around a flat background be denoted by $\hat{\mathcal{L}}_I$. This operator contains in addition to NSNS fluctuations such as $h_{\mu\nu}(x)$, RR fluctuations such as $F_3 = dC_2$. The equations of motion follow from the physical state condition,

$$Q\hat{\mathcal{L}}_I|\Omega\rangle = 0. \quad (1.1)$$

where the state vector above is the ground state corresponding to the spherical world sheet. The above equation translates into linearized equations of motion and Bianchi identities such as

$$\nabla^2 h_{\mu\nu} = 0, \quad d^*F = 0, \quad dF = 0 \dots \quad (1.2)$$

Now we need to allow for the possibility of the world sheet opening up giving the disc topology. The corresponding boundary state $|B\rangle$ was constructed in the nine-brane case in [7] and for the general D-brane in [8] and [10]. While this state is BRST invariant by itself, in order to consider it as a vertex operator on the sphere on the same footing as the other terms one needs a propagator Π , and the total state one should consider is

$$|\Psi\rangle = \hat{\mathcal{L}}_I |\Omega\rangle + \Pi |B\rangle. \quad (1.3)$$

The BRST invariance of this state ($Q|\Psi\rangle = 0$) then leads to modified field equations such as

$$\nabla^2 h = g T_{\{\mu\nu\}} \delta^{9-p}(x_\perp) [\det(1 + \mathcal{F})]^{1/2}, \quad d^* F_5 = i \frac{g}{2} * J, \dots \quad (1.4)$$

In the above $\mathcal{F} = F - B$ where F is the gauge field strength of the open string gauge field and B is the two form NSNS field, T is a certain ($\mathcal{F}$ dependent) matrix in space time, and J is a delta functional current with support at the location of the D-brane.

These arguments admittedly yield only weak field equations in the weak string coupling limit. However they have a unique non-linear generalization. This follows from the fact that the only generally covariant action which will yield the left hand sides of (1.4) is the string effective action (in our case for type IIA or IIB). Similarly the source action turns out to be the DBI action. One is thus led to the conclusion that the effective theory below the string scale is given by the sum of the supergravity (including RR fields) action and the DBI action. i.e.

$$S[\Phi, l] + S_{DBI}[\Phi_l; A], \quad (1.5)$$

where Φ stands for the collection of supergravity fields and Φ_l is the boundary value of these fields at the location (denoted by l) of the D-branes. By A we mean both the gauge field on the brane and the transverse components of the 10d gauge field which represent fluctuations in the location of the brane. To make the discussion precise we will concentrate on the example of the D3 brane and its near horizon geometry[3]. In this case the DBI action becomes the super Yang Mills action and the geometry becomes that of $AdS_5 \times S_5$, but we expect the arguments below to be of more general validity.




Figure 1: The location of the brane in the AdS space. The dashed region is replaced by the brane.

The geometry is given by the $AdS_5 \times S_5$ metric (with Euclidean signature)

$$ds^2 = \frac{R^2}{z^2}(dt^2 + dx_i^2 + dz^2) + R^2 d\Omega_5^2. \quad (1.6)$$

where $R = 4\pi(gN)^{\frac{1}{4}}\sqrt{\alpha'}$. The boundary of this geometry may be taken to be a flat space at $z = 0$ and a point (which for convenience we will call the horizon) at $z = \infty$. Let us discuss quantum field theory in this background with the action (1.5). As in [4] we will only consider fluctuations that are constant on the S_5 so we effectively have a five dimensional theory. The brane is located at $z = l$ and is supposed to give an effective description in the region $l > z \geq 0$ of the region ($z > l$) behind the “stretched horizon” at $z = l$ (see figure). The supergravity action in this region is,

$$S[\Phi, l] = \int_0^l dz \int d^4x L[\Phi]. \quad (1.7)$$

Now consider the quantum effective action (at its minimum) for fields which satisfy the boundary condition $\Phi = \Phi_0$ at $z = 0$. (This quantum field theory is effective at scales below the string scale).

$$e^{-\Gamma[\Phi_0]} = \int [d\Phi]_l |_{\Phi_0} [dA] e^{-S[\Phi, l] - S_{DBI}[\Phi_l, A, l]}. \quad (1.8)$$

We have defined the measure above as $[d\Phi]_l = \prod_0^l d\Phi(z)$. Now the crucial assumption is that Γ is independent of l the position of the brane. It should be stressed that we do not have a justification for this assumption beyond the fact that for minimally coupled scalars the wave function is independent of l . In fact the situation may be more complicated for non-minimal fields [11]. While this does not necessarily make the conjecture invalid it highlights the fact that it is far from being proven.

Thus we may equate the right hand side evaluated at $l = \infty$ to its value at $l = 0$. i.e.

$$\int [d\Phi]_{|\Phi_0} [dA] e^{-S[\Phi, \infty] - S_{DBI}[\Phi_\infty, A]} = \int [dA] e^{-S_{DBI}[\Phi_0, A]} \quad (1.9)$$

Where we have used the fact (see (1.7)) that the supergravity action vanishes for $l = 0$. At scales well below the Planck scale we may approximate the supergravity functional integral by its classical value Φ^c (which satisfies the same boundary conditions as Φ) so that we have the equation,

$$S[\Phi^c, \infty] = \int_0^\infty dz \int d^4x L[\Phi^c] = W_{DBI}[\Phi_0] - W_{DBI}[\Phi_\infty]. \quad (1.10)$$

Here the generating functional for connected correlation functions in the gauge theory W_{DBI} is defined by

$$W_{DBI}[\Phi_l] = -\log \int [dA] e^{-S_{DBI}[\Phi_l, A, l]}. \quad (1.11)$$

Apart from an irrelevant constant (the second term on the right hand side which is independent of Φ_0) the equation (1.10) is the ansatz of [4] and [5].

2 Large N limit of the DBI action

Let us discuss now the precise conditions under which supergravity is valid with the configuration of the D-branes in question being again N coincident D3-branes. The supergravity solution for the ground state of the 3-brane theory is the extremal solution,

$$ds^2 = f^{-1/2}(-dt^2 + \sum_{i=1}^3 dx_i^2) + f^{1/2}(dr^2 + r^2 d\Omega_5^2). \quad (2.1)$$

with

$$f = 1 + \frac{R^4}{r^4}, \quad R \equiv \sqrt{(4\pi gN)^{1/2}\alpha'}. \quad (2.2)$$

We also have for the Ramond-Ramond field strength five-form $F_5 \sim R\Omega_5$ where Ω_5 is the volume of the unit five-sphere and the dilaton is a constant.

The corresponding non-extremal solution for the metric, related to some excited state (thermal state) of the brane theory, is

$$ds^2 = f^{-1/2}(- (1 - \frac{r_0^4}{r^4}) dt^2 + \sum_{i=1}^3 dx_i^2) + f^{1/2}((1 - \frac{r_0^4}{r^4})^{-1} dr^2 + r^2 d\Omega_5^2). \quad (2.3)$$

Here r_0 is the position of the horizon and is also a non-extremality parameter. These solutions are valid up to the horizon provided that the curvature at the horizon in string units is small, i.e. $\alpha'\mathcal{R} \sim (gN)^{-1/2} \ll 1$.

In the regime $r \ll R$ the geometry of the extremal solution is given by dropping the ones in (2.1), (2.2), i.e.

$$ds^2 = \frac{r^2}{R^2}(-dt^2 + \sum_{i=1}^3 dx_i^2) + \frac{R^2}{r^2} dr^2 + R^2 d\Omega_5^2. \quad (2.4)$$

which is the by now well-known $AdS_5 \times S_5$ geometry [3]. Similarly the near horizon geometry of the non-extremal solutions may be obtained by considering the region $r \sim r_0 \ll R$. This gives

$$ds^2 = \frac{r^2}{R^2}(- (1 - \frac{r_0^4}{r^4}) dt^2 + \sum_{i=1}^3 dx_i^2) + \frac{R^2}{r^2} (1 - \frac{r_0^4}{r^4})^{-1} dr^2 + R^2 d\Omega_5^2. \quad (2.5)$$

From these solutions it seems that the proper decoupling limit to consider is $g, \alpha' \rightarrow 0$ with R fixed. We need to send $g \rightarrow 0$ to suppress loop contributions, and we are taking a large N limit with $gN = \frac{R^2}{\alpha'}$ also becoming large to suppress the corrections to supergravity. This is of course very similar to the limit taken in [3] but here the scale which defines the supergravity solution is held fixed (rather than going to zero as in [3]).

Now we need to carry out the same ('t Hooft scaling) limit of the DBI action for D3-branes. As we will see below in this limit all the terms of this action are retained.

The DBI action for N D3-branes (ignoring commutator terms, derivatives of F etc., and using a flat metric and ignoring the non-constant part of the dilaton for simplicity)

is,

$$I = \frac{1}{(2\pi)^3(\alpha')^2 g} \int \text{tr}(\det[\eta_{\mu\nu} + (2\pi\alpha')F_{\mu\nu}])^{1/2}. \quad (2.6)$$

In the above the trace is over $U(N)$, g is the string coupling. Also for simplicity of notation we ignore the distinction between the gauge fields in the brane directions and the transverse fields which are supposed to be the moduli that are related to fluctuations in the positions of the branes. Let us expand the integrand schematically in powers of F^2 . Ignoring the first term (which is proportional to the brane volume) we have,

$$I \sim \frac{N}{gN} \int [\text{tr}F^2 + \alpha'^2 \text{tr}F^4 + \dots + (\alpha')^{2n-2} \text{tr}F^{2n} + \dots]. \quad (2.7)$$

In the above $\text{tr}F^4$ for example stands for a linear combination of $\text{tr}F^4$ and $(\text{tr}F^2)^2$ and we have multiplied and divided the action by N to facilitate the passage to 't Hooft scaling. Note that the Yang-Mills coupling is given by $g_{YM}^2 \sim g$.

Now to get 't Hooft scaling we need to have just a factor of N outside the action as one takes the large N limit. Thus we need to rescale the gauge field by writing,

$$A = \sqrt{gN}\tilde{A}; \quad F = \sqrt{gN}\tilde{F}; \quad \tilde{F} = d\tilde{A} + \sqrt{gN}\tilde{A}^2. \quad (2.8)$$

Then we have,

$$\begin{aligned} I &\sim N \int [\text{tr}\tilde{F}^2 + (\sqrt{gN}\alpha')^2 \text{tr}\tilde{F}^4 + \dots] \\ &\sim N \int [\text{tr}\tilde{F}^2 + R^4 \text{tr}\tilde{F}^4 + \dots + R^{4n-4} \text{tr}\tilde{F}^{2n} + \dots]. \end{aligned} \quad (2.9)$$

where we have identified $R \sim (gN)^{\frac{1}{4}}\sqrt{\alpha'}$ as the scale of the dual supergravity solutions introduced in (2.2). Now since this is the scale which (in effect) should be fixed as one takes the limit $\alpha' \rightarrow 0$ all the terms in the DBI action will contribute.

The above means that we have an infinite number of non-renormalizable terms and the field theory description must break down for energies $E \geq O(R^{-1})$. The important point is that as long as R^{-1} is much less than the Plank scale (which is becoming very large in our limit) there is a regime $E \simeq R^{-1}$ in which higher order terms in the DBI action need to be considered on the same footing as the classical supergravity action. In the infra-red limit $E \ll R^{-1}$ however the theory flows to a point on the conformal fixed

line defined by setting all but the F^2 coupling to zero i.e. to the $\mathcal{N} = 4$ super Yang-Mills theory. This large N low energy limit is in fact equivalent to the ‘double scaling’ limit of [2].

Now in the DBI action, in addition to the gauge fields that we have written down explicitly, there are also the moduli which come from the components of the ten dimensional gauge field that are transverse to the D-brane. Let us call them X . A non-zero expectation value for these can be interpreted as the distance between a probe brane and the rest and should correspond to the distance r in the supergravity solution. This is the same sort of relation that one got in the matrix model. Specifically we expect $r = R^2 X$. The point is that the low energy limit in the gauge theory must correspond to the region $r \sim r_0 \ll R$ in the supergravity solution. In this case one expects the correspondence described in [3] between the the conformal theory on the brane (and its excited states) and the Anti de Sitter space supergravity (2.4) (and its excited version (2.5)). Now as pointed out in [4],[5] the Kaluza Klein modes on the S_5 in (2.4) are of order R^{-1} and these masses agree with the dimensions of chiral operators in the gauge theory. This agreement will of course be preserved by non-renormalization theorems and will not be affected by the DBI terms. On the other hand it is not clear in what sense all of string theory is kept since the string states have masses $O(gN)$ that are becoming large in the supergravity limit.

The above considerations will not affect the agreement of the low energy absorption cross section calculated in the weak coupling gauge theory with the supergravity calculation, [2] since these are protected by non-renormalization theorems, and will continue to agree as one continues from strong to weak coupling keeping the energy low. On the other hand the DBI terms will be seen in the intermediate region as pointed out by [11]. (Of course in (2.6) above one should now keep perturbations around the back ground metric as well as the non-constant part of the dilaton). Our argument shows that even though on the supergravity side only the leading contribution is kept, on the gauge theory side there are contributions (for energies that are low compared to the Planck scale but of the order of R^{-1}) coming from the higher order terms in the DBI action.

3 Radiation from Black Holes in the Dilute Gas Approximation

The ansatz of [3],[4] and [5] is also relevant to the understanding of radiation from brane configurations ¹. The main unresolved issue here (apart from the lack of understanding of Schwarzschild black holes) is the fact that a weak coupling calculation (from say an effective string theory) in flat space gives the correct radiation rate of the corresponding black hole [17],[18]. On the one hand there is a calculation of an S-matrix in tree approximation for the annihilation of modes living on the brane to produce a bulk field which propagates out to infinity. This is a flat space calculation. On the other hand there is the supergravity calculation where one calculates the absorption cross section for these bulk fields in the near horizon geometry. The agreement of the entropy of the extremal black holes calculated in these two ways (i.e. from the effective weakly coupled string picture and the supergravity (Gibbons-Hawking) calculation) was explained by arguing that in this case we are counting the degeneracy of BPS states and therefore this should remain constant as we turn up the coupling. However in this non-extremal, non-BPS, situation this argument will clearly not work. ²

At this point we should discuss the question of where the brane is to be located. As we've argued earlier the physics is independent of this location but some locations may be more convenient than others depending on the particular questions one wishes to address. The standard picture is that the brane is a source for the supergravity solution that corresponds to it. One would then place it at the singularity (which it replaces) or at the horizon and think of it as replacing the configuration behind the horizon. In this case one should not use a flat metric (and constant dilaton etc) but one must consistently use the metric etc. coming from the supergravity solution at this point.³ In this case

¹See [1] for reviews.

²Actually even in the BPS case it is still a puzzle why the counting gives exactly the area entropy of Bekenstein and Hawking.

³Precisely this observation was used recently to resolve an apparent paradox involving 'decoupled' NS five-brane theories [12].

the flat space calculations that are done in order to find the entropy and radiation rates are quite mysterious. One needs to make an argument as to why as one increases the coupling to go from flat space to a black hole configuration, the entropy and the rates remain the same. However the picture that emerges from the discussion in [4],[5] is that the brane (or conformal field theory) is at the boundary of the Anti de Sitter space and thus there is no need for any extrapolation. It seems that this idea can be of relevance for understanding the microscopic calculations of entropy and gray body factors for five and four dimensional black holes. Since it is somewhat simpler we will concentrate on the five dimensional black hole but the argument can be easily extended to the four dimensional case.

The microscopic configuration (D-brane gauge theory) that gives the five dimensional black hole is one where (in type IIB) D-strings carrying momentum are bound to D5-branes. The corresponding supergravity solution (with topology $R^2 \times S^3 \times S^1 \times T^4$ [13] is given (in the string frame) by

$$\begin{aligned}
ds^2 &= f_1^{-\frac{1}{2}} f_5^{-\frac{1}{2}} [-dt^2 + dx_5^2 + \frac{r_0^2}{r^2} (\cosh \sigma dt + \sinh \sigma dx_5)^2 + f_1 dx_i dx^i] \\
&\quad f_1^{\frac{1}{2}} f_5^{\frac{1}{2}} [(1 - \frac{r_0^2}{r^2})^{-1} dr^2 + r^2 d\Omega_3^2] \\
e^{-2\phi} &= g^2 f_1^{-1} f_5 \\
H &= 2r_5^2 \epsilon_3 + 2r_1^2 e^{-2\phi} *_6 \epsilon_3
\end{aligned} \tag{3.1}$$

In the above

$$f_1 = 1 + \frac{r_1^2}{r^2}, \quad f_5 = 1 + \frac{r_5^2}{r^2}, \quad r_1^2 = \frac{gQ_1 \alpha'^3}{V}, \quad r_5^2 = gQ_5 \alpha', \quad r_0^2 \sinh 2\sigma = 2 \frac{g^2 n \alpha'^4}{L^2 V}. \tag{3.2}$$

where V is the volume (divided by $(2\pi)^4$) of the four torus around which the five brane is wrapped, Q_1, Q_5, n are integers giving the numbers of each type of brane and the momentum quantum (in units of $\frac{1}{L}$) and L is the radius of S_1 .

Now as in the discussion after (2.4) the relevant decoupling limit that should be taken here to suppress closed string loop corrections and higher order terms in the α' expansion is

$$g \rightarrow 0, \quad \alpha' \rightarrow 0; \quad gQ_1, gQ_5 \rightarrow \infty, \quad r_1, r_5, r_n \equiv r_0 \sinh \sigma \text{ fixed}, \quad V \sim \alpha'^2. \tag{3.3}$$

Now low energies in the D-brane gauge theory corresponds to the region

$$r, r_0, r_n << r_1, r_5. \quad (3.4)$$

Note that this is equivalent to the so-called dilute gas region studied in [13]. In this regime the metric is

$$\begin{aligned} ds^2 &= \frac{r^2}{R^2}(-dt^2 + dx_5^2) + \frac{r_0^2}{R^2}(\cosh \sigma + \sinh \sigma dx_5)^2 \\ &+ \frac{R^2}{r^2}\left(1 - \frac{r_0^2}{r^2}\right)^{-1}dr^2 + R^2 d\Omega_3^2 + \frac{r_1}{r_5}dx_i^2 \end{aligned} \quad (3.5)$$

where $R^2 = r_1 r_5$. The first three terms which give a space whose coordinates are t, x_5, r is actually the metric of the BTZ black hole [14] in 3 dimensions. To see this⁴ we make the coordinate transformation [16],

$$r^2 + r^2 \sinh^2 \sigma = \rho^2. \quad (3.6)$$

Then we find that the metric becomes,

$$ds^2 = ds_{BTZ}^2 + R^2 d\Omega_3^2 + \frac{r_1}{r_5} dx_i^2. \quad (3.7)$$

where,

$$ds_{BTZ}^2 = -\frac{(\rho^2 - \rho_+^2)(\rho^2 - \rho_-^2)}{\rho^2 R^2} dt^2 + \rho^2 \left(d\phi - \frac{J}{2\rho^2} dt\right)^2 + \frac{\rho^2 R^2}{\rho^2 - \rho_+^2)(\rho^2 - \rho_-^2)} d\rho^2. \quad (3.8)$$

where $M = \frac{\rho_+^2 + \rho_-^2}{R^2}$ and $J = \frac{2\rho_+ r_-}{R}$ are the mass and angular momentum of the BTZ black hole, $\rho_+ = r_0 \cosh \sigma$ $\rho_- = r_0 \sinh \sigma$, and the angular coordinate⁵ $\phi = \frac{x_5}{R}$

It should also be noted that in this regime the dilaton is constant $e^{-2\phi} = g^2 \frac{r_5^2}{r_1^2}$ and that there is a constant Ramond Ramond flux.⁶

⁴The relation of the *extremal* five dimensional black hole to the extremal BTZ black hole was first observed in [15]. In [16] an argument relating the non-extremal black hole to the BTZ black hole was given. Our discussion is an adaptation of the latter.

⁵Note that while R is not necessarily equal to L we would still require that it be integral multiple of L .

⁶Note that the BTZ black hole is (locally) conformally flat. This follows from the fact that locally the metric is diffeomorphic to AdS which is conformally flat.

The near horizon back ground that we have obtained is actually an exact solution of the string background field equations. This is because the BTZ black hole has been shown to be (an orbifold of) the $SL(2,R)$ WZW theory [22] and of course so are S_3 (which is a $SU(2)$ WZW theory and T_4 .⁷The conclusion is that when the parameters of the brane configuration are in the dilute gas region there is a sigma model CFT description in the near horizon region as well as in the asymptotically flat region. This means that one need not confine the discussion to $gN \gg 1$ and perhaps this explains why the effective string calculations agree with supergravity.

This brings us to the question of where the brane (CFT) should be located in relation to the supergravity solution. The usual picture has been to consider the string (D-brane) picture as a model for Hawking radiation in flat space with the black hole being replaced by the collection of D-branes. This is expected to be a weak coupling description with the space being flat. The agreement of the entropy with the geometrical calculation was in that case attributed to supersymmetry. The agreement beyond extremal BPS limit for radiation rates including grey body factors in the dilute gas approximation remained a mystery. In the light of recent work [2][3], [4], [5] and the above considerations a somewhat different interpretation which does not rely on supersymmetry appears to emerge. The point is that the geometry is fixed to be the black hole (or rather string) metric even in the limit $g \rightarrow 0$ (with r_1, r_5 fixed). So while both closed string loops and α' corrections are suppressed a non-trivial background is present. Then one interprets the string action as an effective boundary action which represents the black hole configuration. It should be emphasized that this effective string is in fact a closed string and the fact that the open string coupling gN is large is irrelevant to the calculation. Of course we have made the assumption that the functional integral over the gauge field even in the non-abelian case reduces to a effective string action.

This picture is in accord with the recent observations that the entropy of these black

⁷In fact in this case one can avoid the question of RR field backgrounds by taking the S-dual configuration of NS5-F1 and momentum. It is not clear that this is an advantage though since one still has a magnetic NS background.

holes can be obtained from a boundary conformal field theory [20]. What we are suggesting here is that this picture may give also a natural explanation as to why the emission rates and gray body factors agree with the supergravity calculation.

Thus one may argue as follows. Far from the black hole $r \gg R$ we have flat space. Near the black hole $r \ll R$ we have $AdS_3 \times S_3 \times T_4$. Now the gray body factor calculation [18] depended on solving the Klein-Gordon equation in just these two regions and matching the solutions. Thus one could model the relevant physics by replacing the original geometry by $AdS_3 \times S_3 \times T_4$ and a boundary with a flat metric as in the version of the ansatz for $AdS_5 \times S_5$ discussed in [5]. Thus one expects the process taking place in the bulk to be mirrored by the conformal field theory on the boundary which is in this case an effective string theory. This picture is thus an alternative to the correspondence principle [21] as a way of explaining the agreement between the microscopic calculations and the supergravity calculations of black hole thermodynamics.

On the flat space boundary one has the D5-D1-momentum configuration and the relevant limit is given in (3.3). The important point is that by integrating over the gauge field we get an effective (closed) string in a manner analogous to the way in which effective (closed) (p,q) strings were obtained in ([9],[10]). Thus we expect the boundary gauge theory functional integral

$$\int [dX][dA] e^{iS_{DBI}[\phi^0, A, X]} = \int [dX] e^{iI[\phi^0, X]}. \quad (3.9)$$

where the action in the expression on the right hand side is an effective string action⁸ which we may write schematically as

$$I \sim \frac{T}{2} \int \partial X^\mu \partial X^\nu (\eta_{\mu\nu} + \kappa h_{\mu\nu}). \quad (3.10)$$

where $\kappa \sim g\alpha'^4$. The h field is of course (the boundary value of) an external field coming from the supergravity solution (generically called Φ earlier) and according to the ansatz in [4], [5], one is to regard this as a classical source. But what we have above is precisely the effective string used in [17] to calculate the Hawking radiation rate. Thus the ansatz

⁸A suggestion for deriving this from first principles has been given in [22].

of [4],[5] appears to give an explanation of the agreement between the string calculation and the supergravity calculation. Presumably such arguments can be extended to explain the agreement of the grey body factors [18] as well.

4 Discussion

To conclude let us summarize our results. We have shown that the ansatz of [4] and [5] is equivalent to the assumption that the quantum effective action for the low energy field theory is independent of the location of the brane. Then we showed that the scaling limit that keeps the full supergravity D-brane solutions corresponds to having the full DBI action on the gauge theory side. One is thus led to a relation between the full DBI action with an effective cutoff $R^{-1} < M_p$; $R \sim (gN)^{1/4} \sqrt{\alpha'}$ and supergravity, but in the low energy limit one could argue for a correspondence between for instance the N=4 Yang-Mills theory and supergravity on $AdS_5 \times S_5$. Next we argued that this ansatz provided justification for the effective string model [17] calculations which agreed with absorption rates of (and emission rates from) certain black holes. The argument provides a rationale for treating the effective string as if it were living in flat space and for the non-renormalization of the rates.

5 Acknowledgements:

I wish to thank Igor Klebanov and Edward Witten for useful comments. I would also like to thank Edward Witten for hospitality at the Institute for Advanced Study and the Council on Research and Creative Work of the University of Colorado for the award of a Faculty Fellowship. This work is partially supported by the Department of Energy contract No. DE-FG02-91-ER-40672.

References

- [1] D. Youm, “Black Holes and Solitons in String Theory” hep-th/9710046; A. Peet, “The Bekenstein Formula and String Theory (N-brane Theory)” hep-th/9712253.
- [2] I. Klebanov, *Nucl. Phys.* **B496** (1997) 231; I. Klebanov and S. Gubser, *Phys. Lett.* **B413** (1997) 41, hep-th/9708005.
- [3] J. Maldacena, “The Large N limit of Superconformal Field Theories And Supergravity”, hep-th/9711200.
- [4] S. Gubser, I. Klebanov and A. Polyakov, “Gauge Theory Correlators from Non-Critical String Theory”, hep-th/9802109.
- [5] E. Witten, “Anti-de Sitter Space And Holography”. hep-th/9802150.
- [6] J. Polchinski, “TASI Lectures on D-Branes”, TASI 1996 World Scientific (1997). hep-th/9611050
- [7] C. Callan, C. Lovelace, C. Nappi, S. Yost *Nucl. Phys.* **B308** 221 (1988).
- [8] M. Li, *Nucl. Phys.* **B460** 351 (1996) hep-th/9510161.
- [9] C. Schmidhuber, *Nucl. Phys.* **B467** 146 (1996), hep-th/9601003.
- [10] S. de Alwis and K. Sato, *Phys. Rev.* **D53** 7187 (1996) hep-th/9601167.
- [11] S. Gubser, A. Hashimoto, I. Klebanov, M. Krasnitz, “Scalar Absorption and the Breaking of the World Volume Conformal Invariance” hep-th/9803023; A. Rajaraman, “Two-Form Fields and the Gauge Theory Description of Black Holes”, hep-th/9803082.
- [12] S. de Alwis “Radiation of RR states from NS Fivebranes” hep-th/9801162.
- [13] G. Horowitz and A. Strominger, *Phys. Rev. Lett.* **77** (1996) 2368, hep-th/9602051.
- [14] M. Banados, C. Teitelboim and J. Zanelli, *Phys. Rev. Lett.* **69** (1992) 1849, hep-th/9204099.

- [15] S. Hyun, “U-duality between Three and Higher Dimensional Black Holes” hep-th/9704005.
- [16] K. Sfetsos and K. Skenderis, “Microscopic derivation of the Bekenstein-Hawking entropy formula for non-extremal black holes”, hep-th/971113.
- [17] S. Das and S. Mathur, *Nucl. Phys.* **B482** (1996) 153, hep-th/9607149.
- [18] J. Maldacena and A. Strominger, *Phys. Rev.* **D55** (1997) 861, hep-th/9609026.
- [19] G. Horowitz and D. Welch, *Phys. Rev. Lett.* **71** (1993) 328, hep-th/9302126; N. Kaloper, *Phys. Rev.* **D48** (1993) 2598, hep-th/9303007; A. Ali and A. Kumar, *Mod. Phys. Lett.* **A8** (1993) 2045, hep-th/9303032.
- [20] A. Strominger, “Black Hole Entropy from Near-Horizon Microstates” hep-th/9712251; V. Balasubramaniam and F. Larsen, “Near Horizon Geometry and Black Holes in Four Dimensions”, hep-th/9802198.
- [21] G. Horowitz and J. Polchinski, “Correspondence Principle for Black Holes and Strings” hep-th/9612146.
- [22] S. Hassan and S. Wadia, “Gauge Theory Description of D-brane Black Holes: Emergence of the Effective SCFT and Hawking Radiation”, hep-th/9712213.